

VERTICAL DISTRIBUTIONS OF TEMPERATURE, SALINITY  
AND GEOSTROPHIC FLOW ALONG 155°E AND 170°W  
IN THE SOUTHERN OCEAN IN DECEMBER 1968-  
FEBRUARY 1969 (EXTENDED ABSTRACT)

Mikio NAGANOBU

*Tokyo University of Fisheries, 5-7, Konan 4-chome, Minato-ku, Tokyo 108*

There are large differences in macro-scale oceanic environments among sectors of Atlantic, Indian and Pacific in the Southern Ocean. The HAKUHO MARU worked on oceanographic sections along 155°E and 170°W in the Pacific sector of the Southern Ocean in December 1968–February 1969 (Ocean Research Institute, University of Tokyo (1970): Oceanographic Data of KH-68-4 “Southern Cross Cruise” of the HAKUHO MARU) (Fig. 1). This paper describes vertical distributions of temperature, salinity and geostrophic flow, almost to the sea bottom, from 35°S to near the Continental shelf of Antarctica along 155°E and 170°W.

Along 155°E: The Antarctic Surface Water occupies between 63 and 59°S. The temperature minimum was  $-0.88^{\circ}\text{C}$  at 53 m in the Antarctic Surface Water. The temperature maximum was more than  $1.75^{\circ}\text{C}$  at the layer between 200 and 300 m in the Warm Deep Water. There was a sharp drop in surface temperature from  $7.8^{\circ}\text{C}$  at 54°S to  $1.3^{\circ}\text{C}$  at 62°S. Between 200 and 1000 m, there was also a sharp drop of temperature from  $8.0^{\circ}\text{C}$  at 51°S to  $2.5^{\circ}\text{C}$  at 56°S. In the area having the sharp drop of temperature, a salinity minimum layer of 34.50‰ and under drops from the surface to 1000 m. On the other hand, a salinity maximum layer of 34.72‰ and up rises from 3000 m at 38°S to 500 m at 63°S. Geostrophic flow was calculated referring to the near bottom level. Along 155°E, the water generally flows eastward with a maximum speed of 14.9 cm/s at the surface between 51 and 57°S (Fig. 2). The geostrophic volume transport through 155°E was  $163 \times 10^6 \text{ m}^3/\text{s}$  (eastward flow) between 38 and 63°S.

Along 170°W: The Antarctic Surface Water occupies between 70 and 63°S. The temperature minimum was  $-1.78^{\circ}\text{C}$  at 44 m in the Antarctic Surface Water. The temperature maximum was more than  $1.25^{\circ}\text{C}$  at the layer between 200 and 500 m in the Warm Deep Water. A salinity minimum layer of 34.50‰ and under drops from the surface at 59°S to 1000 m at 38°S. On the other hand, a salinity maximum layer of 34.72‰ and up rises from 3500 m at 38°S to 500 m at 70°S. Geostrophic flow along 170°W generally was eastward with a maximum speed of 12.4 cm/s at the surface between 60 and 62°S (Fig. 3). The geostrophic volume transport through 170°W was  $158 \times 10^6 \text{ m}^3/\text{s}$  (eastward flow) between 38 and 70°S.

*(Received April 28, 1987; Revised manuscript received July 3, 1987)*




Fig. 1. Oceanographic sections along 155°E and 170°W in the Southern Ocean surveyed by the HAKUHO MARU in December 1968–February 1969.




Fig. 2. Vertical distribution of geostrophic flow along 155°E in January–February 1969.




Fig. 3. Vertical distribution of geostrophic flow along 170°W in December 1968–January 1969.